Criteria to Return to Camp After Illness

If a camper or a staff member has symptoms of illness, they should not come to camp.

CRITERIA FOR SYMPTOMS W/ NEGATIVE TEST

Staff cannot return to camp until their symptoms have cleared and they have had no fever for 24 hours without the use of fever-reducing medicines. Campers who need to miss any part of their session due to symptoms of illness will need to reschedule to a different session if possible, due to the nature of our programmatic design. In other words, if the health officer determines that a camper's symptoms are such that they need to leave camp, the camper cannot come back to the same session even if they test negative and meet other criteria to return.

In the case of a confirmed diagnosis of COVID-19, CazPAC uses guidelines from state and local public health agencies as well as CDC recommendations in developing its criteria to return to camp. Persons diagnosed with COVID-19 may exhibit symptoms or may show no detectable symptoms. The criteria for return differ depending on the circumstances.

CRITERIA FOR SYMPTOM-BASED COVID-19

Persons who think or know that they had COVID-19, and had symptoms, may return to camp under the following circumstances:

- At least 10 days have passed since symptoms first appeared; and
- At least 24 hours have passed with no fever, without the use of fever-reducing medications; and
- COVID-19 symptoms have improved (for example, cough, shortness of breath)

CRITERIA FOR TEST-BASED COVID-19

Persons with laboratory-confirmed COVID-19 who had no symptoms will be admitted on site under the following circumstances:

- At least 10 days have passed since the date of their first positive COVID-19 diagnostic test assuming they have not subsequently developed symptoms since their positive test
- If they develop symptoms after testing, the symptom-based criteria override the diagnostic test criteria and begins from the date symptoms present

Returning from Travel

CazPAC requires that all staff members, campers, parents and visitors observe state and local public health guidelines when returning from travel. Those guidelines may include self-quarantining at home and monitoring of symptoms for a period of time, depending on the travel destination, method of travel, and the vaccination status of the group of people traveling.